




Equations, Code, and Salvador Dali for Pi Day

Pi, Part II

MAR 14, 2024




A Jaunty Pi (from [Wikipedia](#))

In [Part I](#) of this essay, I talked about Pi's value and its irrationality. Does that also make Pi Day valuable and irrational? ¹ How about learning more than the 15 digits needed by NASA? Or worthless but rational? Whichever, highly intelligent people are drawn to memorizing digits, just as they are drawn to chess. Think of the celebrated [55-Digit Pi March](#), by Coach Tom and his teenage computers, or how Bruce Dan (see [Wikipedia](#)) recited 144 digits of Pi at a frat party. ² Or think of the many mathematicians over the ages who have devoted time to computing Pi to more and more digits. Even before calculus was invented, Van Ceulen got up to 35 digits in 1610 (see Peter Borwein, "[The Amazing Number Pi](#)" (2000)).

Babylonians	2000? BCE	1	3.125 ($3\frac{1}{8}$)
Egyptians	2000? BCE	1	3.16045 ($4(\frac{8}{9})^2$)
China	1200? BCE	1	3
Bible (1 Kings 7:23)	550? BCE	1	3
Archimedes	250? BCE	3	3.1418 (ave.)
Hon Han Shu	130 AD	1	3.1622 ($=\sqrt{10}$?)
Ptolemy	150	3	3.14166
Chung Hing	250?	1	3.16227 ($\sqrt{10}$)
Wang Fau	250?	1	3.15555 ($\frac{142}{45}$)
Liu Hui	263	5	3.14159
Siddhanta	380	3	3.1416
Tsu Ch'ung Chi	480?	7	3.1415926
Aryabhata	499	4	3.14156
Brahmagupta	640?	1	3.162277 ($=\sqrt{10}$)
Al-Khowarizmi	800	4	3.1416
Fibonacci	1220	3	3.141818
Al-Kashi	1429	14	
Otho	1573	6	3.1415929
Viete	1593	9	3.1415926536 (ave.)
Romanus	1593	15	
Van Ceulen	1596	20	
Van Ceulen	1610	35	

As we talked about last week, Pi is irrational, so there is no end to computations. Unless, that is, you expand your horizons. There are circumstances in which Pi is indeed a rational number, with a finite number of decimal places. In fact, sometimes Pi equals exactly 3.

When I had my students go home and try to measure Pi using a string and a ruler as I described last week, they naturally assumed I meant for them to draw their circles on a flat surface. But suppose they didn't. Suppose they drew their circles on the coach, or on a garbage can. In class, I had Jonathan go find a ball from the Resource Room. Then I drew a circle on it. The definition of a circle drawn on a ball is the same as always: pick a center (the North Pole will do) and go equal distances—on the ball's surface, no cheating by cutting corners and digging tunnels. There are lots of diameter circles you could draw, but go all the way to the equator, which is a circle.



A Sphere (from [Wikipedia](#))

When I did this in class, I asked if Pi would be bigger or smaller than 3.14 if the circle was drawn on a sphere. Smaller, Lyndon said, because the diameter in $\text{Pi} = C/d$ is bigger if you have to go all the way around the curve of the sphere. Then I asked if Pi would be bigger than 2, or less.

I drew a picture, as the first step in solving any problem. I can't draw a good sphere, so I drew a line with a semicircle above it. The line is the flat-circle diameter. The semicircle is the ball-circle diameter, the unknown we're trying to find.

I had them eyeball it and write their estimates on lapboards. 1.1, 2.0, 2.5, 3.0, 2.5. Reasonable guesses, but mostly too big. I asked how to find it more accurately. "Use string" was the good scientific answer, from Emma, I think. I said they could do much better mathematically, with no more information than they had already.


They were stumped. I said I'd give them a couple of minutes. I should have had them discuss it together too. Then Zoe suggested 3.14. "The right idea," I said, "but wrong." even just eyeballing the diagram, 3 is too big.

For the two dimensional circle, $\pi = C/d$. For our equatorial circle, the circumference is the same, C . The diameter, though, is the half-circle going through the north pole. Thus, $\pi = C / (.5C) = 2$.

$\pi = 2$ is the smallest value of π we get on the sphere, and we got it by using the biggest circle that you can draw on a sphere. Consider the Polar Circle, which would have diameter d_2 if you could cut through the globe. I've magnified it in the little picture on the right. The spherical diameter from a_2 to b_2 is still bigger than 2, but not by much. Thus, the π we'd calculate for circle 2 would be close to 3.14. In fact, the closer we got to the North Pole, the closer spherical π would get to 3.14159. . . If we had an infinitesimally small circle at the North Pole, π would take the same value as a flat circle.

How do we get $\pi = 3$? Since we have $\pi = 2$ for the equatorial circle, and $\pi = 3.14$ near the North Pole, we'd have $\pi = 3$ somewhere in between. So sometimes $\pi = 3$. Or, another way to get it would be by keeping the circle the same size as on the equator of this circle, but blowing up the sphere so it has less curvature for a circle of that size— sort of sliding the circle north.

You can think about other drawing shapes on other objects besides flat paper and spheres. Salvador Dali famously drew them on a box, draped from a tree, and on a piece of cloth.



“The Persistence of Memory,” Dali (Wikimedia Commons)

Consider the clock on the box. Suppose we smooth it out so it’s a neat circle instead of being wrinkled. What is the value of Pi?

Well, the circumference is no different than if the watch was flat on the top of the box. Bending the circle over the edge doesn’t change the shape. How about the diameter? Oddly enough, that doesn’t change either. Where the hands meet is the center, and the distance to any edge point is the same as if the watch were flat. So the Dali-Pi is our old friend 3.14.159. . .

Notice that we can lay the watch flat on Dali’s box, with a bend. That’s crucial. You couldn’t do that on a sphere. In fact, you can’t do that just anywhere on a box. You can go over long edges like Dali did, but you can’t paste a circle onto a corner of the box. There, we could draw a circle, as we did on the sphere, by picking a center—the corner itself, for example—and then finding the points equally distant from it. But the value of Pi would depend on the exact point. If I want to finish this while it’s still Pi Day, though, I’d better not go down that rabbit hole. I will note, however, that on the sphere it doesn’t matter where your center is, even though it does depend on the size of your circle. That’s because a sphere has constant curvature. On an ellipsoid or a box it does matter.

Let’s return to flat space, also called Euclidean space. (Did you know you were just doing Non-Euclidean geometry?) But let’s use equations, now. Here’s one by Leibniz, who got it with the help of calculus, which he invented simultaneously with Newton.

$$\pi = 4 \left(1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \dots \right) \quad (\text{Leibniz's formula for } \pi)$$

I won’t try to explain why Leibniz’s formula is true, since that requires calculus, I think (is there any other way, readers?). It also requires trigonometry, which my 7th graders haven’t had yet. But you should be able to see the pattern. I’ll use it later in a computer program that demonstrates it gets close to Pi, though that’s not a proof it gets all the way.

In class, I gave my students a [three-page handout](#) with lots of Pi formulas, most of them too advanced for 7th graders to understand because of the notation. I asked them to pick one each for me to try to explain. Three students picked Euler’s formula and two picked the Normal density formula. Here is [Euler’s formula](#):

$$\pi = \sum_{k=1}^{\infty} k \frac{2^k k!^2}{(2k)!} - 3 \quad (\text{Euler})$$

Euler’s formula is actually written in symbols my 7th graders do learn. It is not that hard to understand, it just needs slow thinking. Suppose $k = 4$. Then $k!$ means $4 \times 3 \times 2 \times 1$, which equals 24. The big Sigma means sum, and it means we add the formula that comes right after it for every k from 1 to infinity. So this formula isn’t as bad as it looks.

My students also picked the formula derived from the [Normal probability density](#), the bell-shaped curve. The formula is:

$$\pi = \frac{1}{2} \left(\int_{-\infty}^{\infty} e^{-x^2/2} dx \right)^2 \text{ (from the Normal density)}$$


This formula is harder to explain, because it is a calculus formula and the special number e (from Euler) an irrational number that equals about 2.7. The fancy symbol is an integral sign, which is a continuous version of the Sigma summation sign in Euler's formula. Euler summed his fraction for $k = 1, 2, 3, \dots$. The integral here sums up for all values of x start from $-\infty$ up to $+\infty$. It doesn't add up to infinity because, like the $1/k$ fractions in Euler's formula, the exponent here gets smaller and smaller.

$$1 = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx \text{ (the Normal density with } \mu = 0, \sigma^2 = 1, \text{ integrated to one)}$$

The normal density can be graphed as in the German diagram below if we also put in a parameter to make the middle number 100 instead of 0, and set the variance (the spread) equal to 15. The area under the bell curve equals 1, as in the integral above, because the probabilities of all possible values of x add up to 1.

$$f(x) = \frac{1}{15\sqrt{2\pi}} e^{-\frac{1}{2} \frac{(x-100)^2}{15}} \text{ (the Normal density with } \mu = 100, \sigma = 15)$$

I use this example because it's one I'm often having to recalculate to figure out which IQ's are which percentiles of the population—for example, 2.2% of people have IQ greater than 130.



The Bell Curve, Centered at 100 (*IHVO Handbook*)

Why should the formula for the Bell Curve also be a formula for Pi? It's a testament to God's glory. The reason involves trigonometry, wave formulas, and something called the Fourier transform and spectral analysis, which I have never got round to learning.

So let's leave formulas now and go to something much less intellectually taxing: computer coding. Below is a Python program that calculates Pi. The program asks the user to tell it how many microseconds of computer time to use (the more you use, the better the estimate of Pi, roughly speaking, or maybe even exactly. Then it prints out two estimates of Pi for you, one using Leibniz's formula and one using Euler's. If you delete some of the pound signs (#), which "comment out" some of the lines, the program will also show its work along the way—its estimates or how much computer time it's used so far.

```
#March 14, 2024. A pi estimator
# Eric Rasmusen, erasmuse61@gmail.com, Python 3

import time
import math

print(f"How many microseconds do you want the computer to use for each
estimate? (e.g. 10, 20000)?")
microseconds = input()
microseconds = int(microseconds)
print("##### ")
print(f"The first method uses Leibniz's formula. ")
start = time.time()

pi = 4
for item in range(1,1000000000):
    if item % 2 == 1: #This means if the remainder from item/2 is 1.
        pi = pi - 4/(2*item+1)
    else: #If the remainder is NOT 1, then item is even.
        pi = pi + 4/(2*item+1)
    end = time.time()
    elapsed = 100000*(end-start)#To measure in microseconds
# print(f"Computer time used was {elapsed:0.2f} microseconds.")
    if elapsed > microseconds:
        break

# print(f"Item is {item} and pi is {pi:0.2f}")

print(f"Pi = {pi:0.15f} using Leibniz.")
print(f"CPU time: { elapsed:0.2f} microseconds.")
print("##### ")

print(f"The second method is Pfouffe's series. ")
start = time.time()

pi = -3
for item in range(1,1000000000):
    pi = pi + item*2**item*(math.factorial(item))**2/math.factorial(2*item)
# print(f"Iteration {item}: pi is {pi:0.2f}.")
```

```

    end = time.time()
    elapsed = 100000*(end-start)#To measure in microseconds
#    print(f"Computer time used was {elapsed:0.2f} microseconds.")
    if elapsed > microseconds:
        break

print(f"Pi = {pi:0.15f} using Euler.")
print(f"CPU time: {elapsed:0.2f} microseconds.")
print("##### ")
print(f"That is all.")
print("##### ")
#I erased the normal distribution method by accident.
#It was fun, but I did need to do an approx of an integral
#Maybe for fun I will try it later.

```

The output looks like this:

```

How many microseconds do you want the computer to use for each estimate? (e.g.
10, 20000)?
16
#####
The first method uses Leibniz's formula.
Pi = 3.146243791226152 using Leibniz.
CPU time: 16.09 microseconds.
#####
The second method is Euler's series.
Pi = 3.141592653471791 using Euler.
CPU time: 16.12 microseconds.
#####
That is all.
#####

```

As you see, Euler's formula works more quickly ("converges faster").

If you'd like to try the program out, go to [Codabrainy.com's free Python page](https://codabrainy.com). Find the box for writing your code in, and clear it out by deleting everything there. Then paste in the program in the first grey box above. Click the sideways triangle at the top to run the program, and you will see the output in the window to the right of the code input window. The program does have a problem, though. It isn't set to print more than about 16 digits of Pi. Later maybe I'll find how Python can be made to calculate and print with more digits.

That is all.

1 How interesting that “value” and “irrational” have so many different meanings! (so many values?) Even “irrational values” has a couple.

2 In my fraternity back in the 1960s we probably had the world champion memorizer of Pi digits, Bruce Dan. He had memorized over 500 digits; alas there was no official record back then.

But 30-odd years later we had a reprise at my home where we all pledged X dollars to our pre-DEI Alma Mater for each digit he could still do, which turned out to be about 144. This event was memorialized here, with the actual recitation of digits starting at about 9:45.

[personal communication, 2024]

8 Comments



Write a comment...



Eric Rasmusen Mar 14 Author

I'll have to remember those till next February.

LIKE REPLY SHARE ...



Amelia Buzzard Writer's Blog(ck) Mar 14

I liked that you gave the 7th graders a sheet of mysteriously complicated looking formulas, instead of just including ones they would understand. It gives them a sense of how much further they could go with math if they wanted to. That sort of pedagogical move should have a name.

 LIKE  REPLY  SHARE

...

6 more comments...

© 2024 Eric Rasmusen · [Privacy](#) · [Terms](#) · [Collection notice](#)
[Substack](#) is the home for great writing